




Envejecimiento saludable y atención primaria

Mikel Izquierdo, PhD.
 Universidad Pública de Navarra
 Departamento de Ciencias de la Salud

Madrid, 15 de Octubre de 2019









WHO ICOPE SCREENING TOOL

Priority conditions associated with declines in intrinsic capacity	Tests	Assess fully if any answer in each domain triggers this
COGNITIVE DECLINE (Chapter 4)	1. Remember three words: flower, door, rice (for example) 2. Orientation in time and space: What is the full date today? Where are you now (home, clinic, etc)? 3. Recalls the three words?	☐ Wrong to either question or does not know ☐ Cannot recall all three words
LIMITED MOBILITY (Chapter 5)	Chair rise test: Rise from chair five times without using arms. Did the person complete five chair rises within 14 seconds?	☐ No
MALNUTRITION (Chapter 6)	1. Weight loss: Have you unintentionally lost more than 3 kg over the last three months? 2. Appetite loss: Have you experienced loss of appetite?	☐ Yes ☐ Yes
VISUAL IMPAIRMENT (Chapter 7)	Do you have any problems with your eyes: difficulties in seeing far, reading, eye diseases or currently under medical treatment (e.g. diabetes, high blood pressure)?	☐ Yes
HEARING LOSS (Chapter 8)	Hears whispers (whisper test) or Screening audiometry result is 35 dB or less or Passes automated app-based digits-in-noise test	☐ Fail
DEPRESSIVE SYMPTOMS (Chapter 9)	Over the past two weeks, have you been bothered by - feeling down, depressed or hopeless? - little interest or pleasure in doing things?	☐ Yes ☐ Yes



ICOPE HANDBOOK APP

Mobile applications will be available to guide health and social care workers on all the steps to undertake, from screening to assessing, to designing a personalized care plan. The app will also produce a printable summary of the results of the assessment and interventions to be included in the care plan in PDF format.

Mejorar la función musculoesquelética, la movilidad y la vitalidad




1. Los **ejercicios multimodales**, que incluyen el **entrenamiento de fuerza con resistencia progresiva** y otras actividades (como el ejercicio del equilibrio y la flexibilidad) y el **entrenamiento aeróbico** están recomendados a las personas mayores que presentan un deterioro de la capacidad física, determinada a través de mediciones de la velocidad de la marcha, la fuerza de prensión y otros parámetros utilizados para conocer el rendimiento físico.
2. A las personas mayores que padecen deterioro se les debe recomendar la **nutrición complementaria por vía oral con asesoramiento dietético**.

La pérdida de fuerza y masa muscular, la reducción de la flexibilidad y los problemas de equilibrio pueden dificultar la movilidad. Los cambios fisiológicos que acompañan al envejecimiento también pueden afectar negativamente al estado nutricional, y este, a su vez, puede repercutir en la vitalidad y la movilidad. Las intervenciones que mejoran la nutrición y fomentan el ejercicio físico, o se integran en los planes de atención y se prestan conjuntamente, pueden reducir, frenar o revertir el deterioro de la capacidad intrínseca.

Prevenir el deterioro cognitivo grave y promover el bienestar psicológico




5. Se pueden ofrecer actividades de **estimulación cognitiva** a las personas mayores que presentan deterioro cognitivo, con independencia de si se les ha diagnosticado o no demencia.
6. Asimismo, se pueden ofrecer **intervenciones psicológicas breves y estructuradas** a otros adultos que experimenten síntomas depresivos, de conformidad con los directrices de intervención nHAP de la OMS. Estas intervenciones deben prestarlas profesionales sanitarios que conozcan bien la atención a la salud mental de las personas mayores. Muy a menudo, una misma persona presenta deterioro cognitivo y dificultades psicológicas, problemas que afectan a su capacidad para llevar a cabo actividades cotidianas (como hacer la compra o manejar las finanzas personales) y a su socialización. Tanto la terapia de estimulación cognitiva (un programa compuesto por actividades centradas en temas diversos) como las intervenciones psicológicas breves son fundamentales para evitar que las personas mayores sufran pérdidas significativas de capacidad mental que dependan de los servicios de asistencia.

Prevenir las caídas




9. Es recomendable revisar los medicamentos que toman las personas expuestas al riesgo de caídas, así como retirar aquellos que sean innecesarios o puedan ser dañinos.
10. Otra intervención recomendada para estas personas son los ejercicios multimodales (equilibrio, fuerza, flexibilidad y entrenamiento funcional).
11. También se han de evitar las fuentes de peligro de caídas. Con este fin, un especialista debe llevar a cabo una evaluación que permita realizar cambios en el hogar para eliminar las fuentes de peligro de caídas presentes en el entorno.
12. Las intervenciones multifactoriales que integran la evaluación y las intervenciones adaptadas a cada individuo también pueden reducir el riesgo y la incidencia de las caídas en las personas mayores.

Las caídas son la principal causa de hospitalización y de muertes relacionadas con traumatismos en las personas mayores. Suelen producirse por una combinación de factores del entorno (alfombras sueltas, desorden en el hogar, mala iluminación, etc.) y factores individuales (anomalías de órganos o sistemas que afectan al control postural). El ejercicio, la fisioterapia, la evaluación y adaptación de las fuentes de peligro en el hogar y, cuando sea necesaria, la retirada de psicofármacos permiten reducir el riesgo de caídas en las personas mayores.

Mantener las funciones sensoriales




3. Es necesario realizar **cribados rutinarios** a las personas mayores en los centros de atención primaria y, si se les encuentran **deficiencias visuales**, prestarles sin demora servicios integrales de atención oftálmica.
4. Además, se debe someter a estas personas a **pruebas para detectar** a tiempo las pérdidas de audición y, si es necesario, suministrarles **audífonos** con el fin de contrarrestarlas.

El envejecimiento causa a menudo disminuciones de la visión y la audición que limitan la movilidad, reducen el compromiso y la participación social y aumentan el riesgo de caídas. Hay medidas sencillas y económicas para hacer frente sin dificultad a los problemas sensoriales, como las lentes correctoras y los audífonos, la cirugía de cataratas y las adaptaciones del entorno.

Hacer frente a problemas asociados a la edad, como la incontinencia urinaria




7. Promover el vaciamiento motivado para manejar la incontinencia urinaria en las personas mayores con deterioro cognitivo.
8. Las mujeres de edad avanzada que presentan incontinencia urinaria (ya sea de urgencia, de esfuerzo o mixta) pueden realizar un **entrenamiento muscular del suelo pélvico**, solo o combinado con técnicas de control vesical y de autovigilancia.

Una tercera parte de las personas mayores presentan incontinencia urinaria (es decir, pérdidas involuntarias de orina), un problema mundial con consecuencias psicosociales como pérdida de autoestima, reducción de la socialización y las relaciones sexuales, y depresión. El entrenamiento muscular del suelo pélvico refuerza los músculos de soporte de la uretra y aumenta su presión de cierre, limitando eficazmente las pérdidas de orina.

Apoyar a los cuidadores




13. Es necesario ofrecer **intervenciones psicológicas, formación y apoyo** a los familiares y otros cuidadores informales de personas mayores que necesitan atención, sobre todo – pero no exclusivamente – cuando estas necesidades son amplias y complejas y cuando los cuidadores están sometidos a una sobrecarga considerable.

Los cuidadores de personas con grandes reducciones de la capacidad funcional están expuestos a un riesgo más elevado de sufrir estrés patológico y depresiones. El estrés y la sobrecarga que puede producir esta actividad tienen un impacto profundo en el estado físico, emocional y económico de las mujeres y de otros cuidadores no remunerados. Es necesario ofrecer a los cuidadores sometidos a este estrés una evaluación de sus necesidades y el acceso a formación y apoyo psicosocial.

Hacia un Envejecimiento Saludable...

Mejorar la función musculoesquelética, la movilidad y la vitalidad



1. Los **ejercicios multimodales**, que incluyen el **entrenamiento de fuerza con resistencia progresiva** y otras actividades (como el ejercicio del equilibrio y la flexibilidad y el entrenamiento aeróbico) están recomendados a las personas mayores que presentan un deterioro de la capacidad física, determinada a través de mediciones de la velocidad de la marcha, la fuerza de prensión y otros parámetros utilizados para conocer el rendimiento físico.
2. A las **personas** mayores que padecen desnutrición se les debe recomendar la **nutrición complementaria por vía oral con asesoramiento dietético**.

La pérdida de fuerza y masa muscular, la reducción de la flexibilidad y los problemas de equilibrio pueden dificultar la movilidad. Los cambios fisiológicos que acompañan al envejecimiento también pueden afectar negativamente al estado nutricional, y este, a su vez, puede repercutir en la vitalidad y la movilidad. Las intervenciones que mejoran la nutrición y fomentan el ejercicio físico, si se integran en los planes de atención y se prestan conjuntamente, pueden reducir, frenar o revertir el deterioro de la capacidad intrínseca.




World Health Organization

Care pathways to improve mobility

5

Locomotor capacity

Care pathways to improve mobility





Mobility is a critical determining factor for healthy ageing. It is important for maintaining autonomy and preventing dependence on care. A person's bodily capacity to move from one place to another is termed locomotor capacity.

Many older people and their families accept losses of locomotor capacity and the associated pain as inevitable. They are not. Indeed, there are effective strategies to improve and maintain mobility in older age.

KEY POINTS

- Limited mobility is common among older people but not inevitable.
- Community-level health-care workers can screen for limited mobility with simple tests.
- A programme of regular exercise, tailored to individual capacities and needs, is the most important approach to improve or maintain locomotor capacity.
- Adapting one's environment and using assistive devices are good ways to maintain mobility despite reduced locomotor capacity.

25



World Health Organization

Care pathways to improve mobility

MANAGE
LIMITED MOBILITY

5.1 MULTIMODAL EXERCISE PROGRAMME

For those with limited mobility, a multimodal exercise programme should be tailored to suit individual capacity and needs.

A multimodal exercise programme for people with limited mobility can include:

- **strength/resistance training**, which requires muscles to work under load, using weights, resistance bands or body-weight exercises such as squats, lunges and sit-to-stand exercises;
- **aerobic/cardiovascular training**, such as fast walking or cycling that increases heart rate until the person is slightly out of breath but can maintain a conversation;
- **balance training**, which challenges the balance system, including static and dynamic exercises; can progress to different surfaces and with eyes open and shut; examples are standing on one leg at a time and walking heel-to-toe in a straight line; and
- **flexibility training**, which improves the extensibility of soft tissues, such as muscle, and the range of joint movement; examples are stretching and other yoga and Pilates exercises.

The Vivifrail project offers a practical guide to developing a tailored exercise programme.

<https://www.vivifrail.com/resources>

Nutrition. Increased protein intake and other nutritional interventions can enhance the benefits of an exercise programme. See Chapter 6 on malnutrition. → 6

Safety of exercise. Before giving advice on exercise or planning an exercise programme, ask about health conditions that would affect the timing or intensity of the activity.

If the person answers yes to any of the following questions, a skilled health professional should develop a tailored exercise programme.

- Have you had chest pain when at rest?
- Have you had a heart attack within the last six months?
- Have you fainted or lost consciousness?
- Have you fallen in the past 12 months?
- Have you broken a bone in the last month?
- Do you get out of breath doing ordinary daily activities at home, such as getting dressed?
- Do you have a joint or muscle disease that limits exercise?
- Has a health-care provider told you to limit exercise?

5

Locomotor capacity

Care pathways to improve mobility

World Health Organization

Care pathways to improve mobility

5

Locomotor capacity

Care pathways to improve mobility

Managing limitations. Where pain limits mobility, pacing physical activity in manageable chunks of time and slowly increasing physical tasks helps to build the body's resilience and manage pain. For people with severely reduced mobility, exercise training is best or watered as a chair can be a starting point. For people with limitations in cognition, such as dementia, a simple and less structured exercise programme may be more suitable.

5.2 SUPPORT FOR SELF-MANAGEMENT

Support for self-management increases adherence to and the benefits of a multimodal exercise programme. People whose SPSS scores are in the range of 10–15 can exercise at home and in the community. People with more severe mobility limitations may need supervision and guidance during exercise.

The WHO mobile health for ageing (mAgging) handbook explains how a mobile phone app can complement health-care professionals' routines and by supporting self-care and self-management.

More information: <http://www.who.int/magging/health-systems/magging>

WHO'S GLOBAL RECOMMENDATIONS ON PHYSICAL ACTIVITY

All older adults can benefit from advice on the physical activity recommended for their age, taking into consideration their health conditions. This box summarizes WHO's global recommendations on physical activity for people aged 60 years and older.

- ✓ Throughout each week, get at least 150 minutes of moderate intensity aerobic physical activity or at least 75 minutes of intensive aerobic activity, or an equivalent combination.
- ✓ Exercise at least 10 minutes at a time.
- ✓ For additional benefits, do 300 minutes of moderate intensity aerobic exercise per week or 150 minutes of intensive aerobic activity per week, or an equivalent combination.
- ✓ Do muscle strengthening activities two days a week or more.
- ✓ If mobility is poor, perform physical activity that enhances balance on three days a week or more.
- ✓ If mobility is poor, perform physical activity that enhances balance on three days a week or more.
- ✓ If you cannot exercise as much as recommended, be as physically active as you can.
- ✓ More information: <http://www.who.int/dietphysicalactivity/physical-activity>

5.3 POLYPHARMACY

Some drugs can impair mobility or interfere with balance per se are sometimes unnecessary or ineffective for a specific person (8). These include, but are not limited to, the following:

- anticholinergics
- benzodiazepines
- neuroleptanalgesic hypnotics
- opioid analgesics
- selective serotonin reuptake inhibitor (SSRI) antidepressants
- antiepileptics
- spasmolytics

Discontinuing unnecessary, ineffective medications as well as medications with a duplicate effect reduces polypharmacy. If the drugs above are used, a medication can be safely stopped, after an appropriate specialist.

5.4 PAIN

Acute pain. Severe pain associated with movement can limit or prevent exercise. It is helpful to ease the severity of pain related to mobility, both to help with designing an exercise programme and for managing the pain. You can use the brief pain inventory: https://www.who.int/pain/assessment_tools/brief-pain-inventory

Manage pain (8). Musculoskeletal conditions that impair mobility often involve persistent pain. A specific biological cause of persistent pain can rarely be found, however. A best practice approach to pain management therefore addresses multiple factors that may be associated with pain – physical factors such as muscle strength, range of movement and endurance, psychological well-being, emotions and sleep. Where pain is a significant barrier to movement and activity, a health professional with specialised knowledge of pain management should develop the pain management plan.

Interventions for pain include:

- self-management → 3.1
- exercises and other physical activity
- medications ranging from paracetamol and non-steroidal anti-inflammatory drugs to gabapentin and opioids
- manual therapy such as massage, joint manipulation and joint mobilisation
- psychological therapy and cognitive behavioural therapy (see Chapter 8 on depressive symptoms) → 8
- acupuncture
- spinal physiotherapy and injections
- radiofrequency denervation.

Some of these interventions can be made available in the community. Others would likely require referral to a specialist facility.

ASSESS & MANAGE
ASSOCIATED CONDITIONS

5

Locomotor capacity

Care pathways to improve mobility


¿De qué manera estamos diseñados para movernos ?



"We are, deep down, still runners, whether or not we declare it by our actions"

Why We Run: A Natural History by Bernd Heinrich (May 7, 2002)

Hemos dejado de movernos y enfermamos



Khan KM, Thompson AM, Blair SN, Sallis JF, Powell KE, Bull FC, Bauman AE.
Sport and exercise as contributors to the health of nations. Lancet. 2012 Jul 7;380(9830):59-64

Ejercicio físico como fuente de salud

Actividad Física y riesgo cardiovascular




Menor mortalidad cardiovascular

(Morris et al. Lancet, 1953. Londres) (Paffenbarger N.Engl.J.Med. 1970)


Relationship Between Ideal Cardiovascular Health and Disability in Older Adults: The Chilean National Health Survey (2009–10)

Antonio García-Hermoso, PhD,* Robinson Ramírez-Vélez, PhD,[†] Rodrigo Ramírez-Campillo, PhD,[‡] and Mikel Izquierdo, PhD[§]




COSTE ECONÓMICO MUNDIAL DE LA INACTIVIDAD FÍSICA

Un estudio analizó los gastos sanitarios de 142 países (93,2% de la población mundial) derivados de cardiopatías, accidentes cerebrovasculares, diabetes tipo II, cáncer de mama y cáncer de colon atribuibles a la inactividad física.



Además de la morbilidad y la mortalidad prematura, la inactividad física es responsable de una enorme **carga económica**.



La **inactividad física** costó a los sistemas de salud mundiales **\$53.800 millones** en 2013.

\$31.200M Sector público
\$12.900M Sector privado
\$9.700M Hogares



Estos datos proporcionan una razón adicional para priorizar la **promoción de actividad física** en todo el mundo.

Ding, D., et al. The Lancet, 2016

fissac.com

Ejercicio, Fármacos y Mortalidad



REVIEWS

Exercise is the Real Polypill

The concept of a "polypill" is receiving growing attention to prevent cardiovascular disease. Yet similar if not overall higher benefits are achievable with regular exercise, a drug-free intervention for which our genome has been shaped over evolution. Compared with drugs, exercise is available at low cost and relatively free of adverse effects. We summarize epidemiological evidence on the preventive/therapeutic benefits of exercise and on the main biological mediators involved.

Carmen Fiuza-Luces,^{1,2}
Nuria Garatachea,³
Nathan A. Berger,⁴ and
Alejandro Lucia^{1,2}

¹Universidad Europea Madrid, Madrid, Spain; ²Instituto de Investigaciones Fisiológicas y de Ciencias de la Salud y del Deporte, Universidad de Zaragoza, Huesca, Spain; and ³Center for Scientific Health and Society, Case Western Reserve University, School of Medicine, Cleveland, Ohio; ⁴alejandro.lucia@eu.edu





FIGURE 2. Summary of the main myokines, their putative effects, and the molecular signals/pathways involved

Exercise benefits for cardiovascular disease go beyond traditional risk factors

Fiuza-Luces C, Santos-Lozano A, Joyner M, Carrera-Bastos P, Zugaza J, Izquierdo M, and Lucia A 2018


nature
REVIEWS CARDIOLOGY





Main beneficial effects of muscle-released myokines on cardiovascular health.

¿ “Envejecimiento” o el estudio de las personas mayores?

Modelo óptimo de envejecimiento



Deficiencia de Ejercicio Físico: La verdadera causa de las enfermedades



Age and sex

REVIEWS

PHYSIOLOGY 32: 152–161, 2017. Published February 22, 2017; doi:10.1152/physiol.00029.2016

Physical Activity, Aging, and Physiological Function

Stephen D. R. Harridge and Norman R. Lazarus

Centre of Human & Aerospace Physiological Sciences, Faculty of Life Sciences & Medicine, King's College London,






FIGURE 1. Depiction of the trajectory of age-related decline in physiological function. **A:** depiction of the trajectory of age-related decline in physiological function (such as $\dot{V}O_{2\max}$ or muscle power) in master athletes, active individuals, and sedentary people. The graph depicts that the age-function relationship is moved upward and to the right as levels of activity increase. This results in a higher age for the same level of physiological function. The graph also depicts the highest values for any age occurring in the master athletes but also a relatively greater loss in $\dot{V}O_{2\max}$ in athletes, associated with a decline in training load. Linear declines in function are shown for illustrative purposes but should not be assumed. **B:** $\dot{V}O_{2\max}$ values taken from Wilson and Tanaka (Ref. 10, men, bottom) and Tanaka and Smith (Ref. 8), women, top) have been allocated to bandwidths of functional equivalence (0–10 $\text{ml} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$). The data for men represent mean values from studies used in the meta-analysis, and for women each data point represents the raw value for a given individual. The graphs separate vigorous exercisers and sedentary people, and show that bandwidths representing different functional equivalence encompass a wide range of ages. Note that the data points plotted are only those that could be discerned from the original publication and that a number of points are overlaid (50).

Norman and Lazarus, 2016 Physiology




¿El ejercicio previene el efecto del envejecimiento?



Valenzuela et al. 2019

COMPREHENSIVE
PHYSIOLOGY

¿El ejercicio previene el efecto del envejecimiento?




Harridge & Saltin (2007) – Adapted from Young (1995)
Encyclopaedia of Gerontology

Journal of Cachexia, Sarcopenia and Muscle (2019)
 Published online in Wiley Online Library (wileyonlinelibrary.com) DOI: 10.1002/jcsm.12432

Effectiveness of a multimodal intervention in functionally impaired older people with type 2 diabetes mellitus



Loredo Rodríguez-Maheú¹, Olga Lasa², Bruno Velho³, Giuseppe Padua⁴, Eva Topinkova⁵, Juan Oliva-Morales⁶, Isabelle Bourde-Marchasson⁷, Michel Cosentino⁸, Kerry Hand⁹, Anding Zou¹⁰, Giovanni Garibani¹¹, Mirko Petrucci¹², Tim C. Hardman¹³, Mark J. Kozak¹⁴, Ivan Badier¹⁵, Gabor Abella¹⁶, Michela Barletta¹⁷, Luc M. Pella-Langlois¹⁸, Sophie C. Requier¹⁹, Riccardo Calvani²⁰, Stefano De Beyer²¹, Alan J. Smeets²² & on behalf of the European MIO-Frail Consortium



- 1 • Exercise Intervention: 16 weeks of resistance training, twice a week, repeated in year 2
- 2 • Nutritional and Educational Intervention: 7 Educational sessions
- 3 • Optimisation of glycaemic and blood pressure control
 HbA1c: 7-8% (9.6-11.6 mmol/L).
 BP <150/90 mmHg.

12 month structured multimodal intervention programme across several clinical settings in different European countries leads to a clinically relevant and cost-effective improvement in the functional status of older frail and pre-frail participants with type 2 diabetes mellitus

Work Programme 2011





lower health care cost per patient vs. usual care of 428.02 € with incremental gain in SPPB score per patient included at the intervention programme vs. usual care during the follow-up period

MIOFrail EUROPEAN COMMISSION / European Research Area / Health & Life sciences

Effect of Exercise Intervention on Functional Decline in Very Elderly Patients During Acute Hospitalization

A Randomized Clinical Trial



Effect of Exercise Intervention on Functional Decline in Very Elderly Patients During Acute Hospitalization

A Randomized Clinical Trial

Table 2. Results of Primary and Secondary End Points by Group^a

Variable ^b	Control Group	Intervention Group	Between-Group Difference (95% CI)	P Value Between Groups
Primary End Point: Change in Functional Capacity				
SPPB score (balance, gait ability, leg strength)	6.2 (4.2 to 8.2)	2.4 (2.1 to 2.7)	2.2 (1.7 to 2.6)	<.001
Barthel index (ADL)	-5.0 (-6.2 to -3.8)	1.9 (0.2 to 3.7)	6.9 (4.4 to 9.3)	<.001
Secondary End Points				
Cognitive decline				
MMSE	0.3 (-0.1 to 0.6)	2.3 (2.7 to 2.0)	1.4 (1.3 to 2.3)	<.001
Orientation (QSO)	0.7 (0.4 to 1.0)	-2.3 (-2.7 to -1.9)	-2.4 (-2.5 to -1.8)	<.001
QSO (Barthel)	-2.2 (-1.8 to 1.0)	11.0 (7.5 to 14.5)	13.2 (8.2 to 18.2)	<.001
Incident delirium (CAM), %	8.3	14.8	OK, 1.0 (0.0 to 4.0)	.12
Readmission (month), %	-0.5 (-2.1 to -1.0)	1.5 (1.1 to 1.8)	2.1 (1.6 to 2.6)	<.001

Abbreviations: ADL, activities of daily living; CAM, Confusion Assessment Method; SPPB, Short Portable Mental Status Questionnaire; QSO, Yesavage Geriatric Depression Scale; MMSE, Mini-Mental State Examination; OK, odds ratio; QSO, quality of life; SPPB, Short Portable Mental Status Questionnaire. ^aAll data, except for CAM, were derived from the final effects model. For each group, data are expressed as change from baseline (admission) to discharge, determined by the linear coefficient (95% CI) of the model. For example, for the SPPB score, 2.2 corresponds to the coefficient estimated from the model. The between-group difference was determined with data on group interaction coefficient. For CAM, data are the proportion of patients with incident delirium during hospitalization.

^bExplanations of variables used are given in the footnotes to Table 1.

Table 3. Results of Secondary End Points Indicative of Adverse Events or Hospitalization

End Point	Control (n = 155)	Intervention (n = 153)	P Value Between Groups
Length of hospital stay, median (IQR), d	8 (4)	8 (4)	.25 ^a
Falls during hospitalization, No./No. (n per group experiencing ≥1 fall)	0/139	4/146 (2.7)	.12 ^a
3-mo Hospital readmission rate (10-person/3-mo), median (IQR)	2.5 (1.8-3.3)	2.4 (1.7-3.2)	.82 ^a
3-mo Mortality, %	9.7	11.9	.62 ^a
Transfer, %			
Home	91.4	92.4	
Institutionalization	1.1	2.2	.55 ^a
Other	7.6	5.4	

Abbreviations: ADL, activities of daily living; IQR, interquartile range.


^aMann-Whitney test.

^bFisher exact test.


^cMid-P value exact test.

^dχ² Test.


A Change in Barthel index from admission to discharge




C Change in Barthel index



B Change in SPPB from admission to discharge



D Change in SPPB





Vivifrail programa de intervención en la fragilidad y riesgo de caídas



PROYECTO VIVIFRAIL

El proyecto Vufval es un programa de Promoción del Ejercicio físico que es referente internacional de intervención comunitaria y hospitalaria para la prevención de la fragilidad y las caídas en las personas mayores. Actualmente está siendo usado por más de 1000 profesionales sanitarios actuando en conjunto sobre la población efectiva de más de 15.000 personas mayores.


La mayoría de novedades en los resultados de fuerza y Endurance que fueron evaluadas durante los estudios indican que la prescripción de ejercicio debe ser



[PROYECTO](#)
[SOCIOS](#)
[EVENTOS-NOTICIAS](#)
[RECURSOS](#)
[CONTACTO](#)

Co-funded by the European Union

 Erasmus+



MATERIALES VIVIFRIL

Vivifral dispone de diversos materiales destinados los profesionales sanitarios deseen intervenir. Las descargas incluyen información sobre la correcta prescripción del ejercicio y Pasaportes que pueden ser utilizados por los pacientes para hacer el seguimiento del programa.

[IR A LA PÁGINA DE RECURSOS](#)



El plan de trabajo incluye la formación presencial o a distancia, y se centrará en:

- A. Profesionales y asistentes de ayuda a domicilio.
- B. Monitores responsables de la gestión de los programas de ejercicio en administraciones públicas, clubes de jubilados y asociaciones.
- C. El personal de centros de salud de atención primaria, secundaria y terciaria.
- D. El personal médico y no médico de residencias de ancianos.

¿TIENES 70 AÑOS O MÁS?

¿TE GUSTARÍA TENER UNA MEJOR CALIDAD DE VIDA?

SIGUE ESTOS PASOS



CONSULTA
CON TU
MÉDICO



HAZ EL TEST
VIVIFRAIL Y
CONSIGUE TU
PASAPORTE



REALIZA EL
PROGRAMA
DE EJERCICIOS



MEJORA TU
CAPACIDAD
FUNCIONAL

¡CONSIGUE EL PASAPORTE!
Y COMPLETA EL PROGRAMA DE EJERCICIOS









Centro de Investigación Biomédica en Red
Fragilidad y Envejecimiento Saludable



NAVARRABIOMED
CENTRO DE INVESTIGACIÓN BIOMÉDICA



With the support
of the European
Commission and the
European Union




aptitude



Interreg
POCTEFA

COMPONENTES DEL TEST VIVIFRAIL Y PROGRAMAS DE EJERCICIO FÍSICO RECOMENDADO



A

DISCAPACIDAD
SPPB 0-3
VIA (AA) < 0.5 m/s

B

FRAGILIDAD
SPPB 4-6
VIA (AA) 0.5 - 0.8 m/s

C

PRE FRAGILIDAD
SPPB 7-8
VIA (AA) 0.8 - 1 m/s

D

BIENESTAR
SPPB 9-12
VIA (AA) > 1 m/s





TEST VIVIFRAIL

El test VIVIFRAIL te permite determinar el tipo de programa físico multicomponente más recomendable a la capacidad funcional y riesgo de caídas de la persona mayor.

El test VIVIFRAIL consiste en valorar la capacidad funcional mediante el SPPB y una batería de 4 pruebas que permiten evaluar el riesgo de caídas. En caso de tener limitaciones para realizar el SPPB, se recomienda medir la velocidad de marcha en 6 metros.

Este test se puede aplicar en un tiempo aproximado de 15 minutos. Se compone de 7 pruebas. Una vez realizadas las pruebas puedes seguir el esquema que se muestra en la página 17 para obtener un resultado.



Centro de Investigación Biomédica en Red
Fragilidad y Envejecimiento Saludable



NAVARRABIOMED
CENTRO DE INVESTIGACIÓN BIOMÉDICA



With the support
of the European
Commission and the
European Union



aptitude



Interreg
POCTEFA


















76

Programa multicomponente de ejercicio físico para el anciano frágil y reducción del riesgo de caídas




Co-funded by the European Union





Descarga nuestra App para realizar los programas de entrenamiento completos.






18

Resources

[Home](#) > [Resources](#)

Resources EN

Materials for professionals responsible for the prescription of a program of physical exercise

In this section you will find all the information and tools necessary for the individualized prescription of a multicomponent program of physical exercise. Download the whole material package through [this link](#), or download only that you need through the links shown below:



Vivifrail Physical Exercise Passport for Professionals

The passport of physical exercise is aimed at professionals and specialists in physical exercise. You can find all the necessary steps for the implementation of the Vivifrail program.

[DOWNLOAD](#)



Exercise wheels by functional level type

Download the specific exercise wheels for each functional level that include the exercises, series and repetitions that should be done every week.

[DOWNLOAD](#)



Vivifrail promotion poster for Exercise Prescription Centers

Download the poster to promote the participation in the Vivifrail program of elderly people who come to centers and consultations. Step by step the way to participate in the exercise program that best suits your initial situation is described.

[DOWNLOAD](#)



Vivifrail Test Poster

Download the poster that explains the components of the Vivifrail test and the recommended exercise programs. If you deploy it in your workplace, it will facilitate the prior assessment process necessary for the recommendation of an individualized program.

[DOWNLOAD](#)






Prescription materials

Passports of Vivifrail Physical Exercise for Participants

Depending on the functional level and risk of falls that each participant has, up to six different types of physical exercise programs can be downloaded. Passports with the "+" symbol are for those with risk of falls.



Passports of Vivifrail Physical Exercise for Participants

Download the entire passport package through this link.

[DOWNLOAD](#)



Passport type A

Download the Passport type A for people with disabilities.

[DOWNLOAD](#)



Passport type B

Download the Passport type B for people with fragility.

[DOWNLOAD](#)



Passport type B+

Download the Passport type B for people with fragility and risk of falls.

[DOWNLOAD](#)



Passport type C

Download the Passport type C for people with pre-fragility.

[DOWNLOAD](#)



Passport type C+

Download the Passport type C+ for people with pre-fragility and risk of falls.

[DOWNLOAD](#)



Passport type D

Download the Passport type D for robust people.

[DOWNLOAD](#)

App and e-book

Download e-book with videos of the exercises and extended information about Vivifrail. You can also install the Vivifrail App that will allow you to digitally track the exercise program.




¿Es Etico NO prescribir actividad física en las personas (a todas las edades)?



- **Ausencia total de** práctica semanal de actividad física (AF) responsable a nivel global de mas de 5 millones de muertes y, \$120 Billones de costo por año
- **Inactividad Física** elemento clave a todas las edades:

"Revertir el Síndrome de Deficiencia de Ejercicio, con especial atención en la mejora de la fuerza muscular"



- Promoción de PA todavía **NO** está incluido en la cartera **de servicios para la prevención y manejo de enfermedades en el entorno de la medicina**
- Aunque puede ser un buen comienzo, la prescripción de AF and ejercicio como medicina **es mucho mas que "usted camine" o "levante la pesa 5 veces"**.

¿Quieres saber más?



mikel.izquierdo@gmail.com



#mikelizquierdo_



#mikelizquierdo_

viviFrail

<http://www.vivifrail.com>



Recent publications: https://www.researchgate.net/profile/Mikel_Izquierdo3/contributions

Publications on PubMed: <http://goo.gl/zaiGw8>

ciberfes
Centro de Investigación Biomédica en Red
Fragilidad y Envejecimiento Saludable

NAVARRABIOMED
CENTRO DE INVESTIGACIÓN BIOMÉDICA

viviFrail



upna
Universidad Pública de Navarra
Unibertsitatea Publikoa

excelencia internacional
ngaiarteko bikaintasuna

Mikel Izquierdo
Universidad Pública de Navarra
Departamento de Ciencias de la Salud